

LADY JANE

AGRIA x CMK2005-709-005







Building on potential

Added value of the variety

Robust plant, tolerant to climatic stress.
Good taste - premium frying quality.
Excellent frying colour throughout the storage period.
Variety with high dry matter content.













Characteristics


Shape of tuber	LO	
Av. Tuber length (Ø 50-60 mm)	9,0 cm	
Maturity (9=early, 4=late)	6,1	
Colour of skin	DY	
Colour of flesh	7,6	
Suitability for cooking	8 C	

Soil adaptability

	Sandy	Light	Heavy
Yield (Relative)	117	120	116
DM (%)	21,9	23,0	22,9
UWW	404	426	426
Tuber size (Ø mm)	51	58	62

Ro1	9	
Ro2/3	3	
Ro5	6	
Pa2	3	
Pa3	1	
wz 1 (D1)	10	
wz 2 (G1)		
wz 6 (O1)	4	
wz 18 (T1)	4	
TRV	6	
Moptop virus	Susceptible	




Climate adaptability

Drought tolerance	Rather tolerant
Secondary growth	7 






Fertiliser requirement

	N	P	K	Mg	B	Ca	Mn
Quantity	-10%	Average	High (+25%)	Average	Average	Average	Average
No. applications	2	1	2	6	2	2	1

Chemical requirement







Foliar late blight	9 
Tuber late blight	Rather tolerant
Early blight	Rather tolerant
Y-virus	8 
Yntn-virus	7 
Leafroll virus	
Black dot	Rather tolerant
Silver scurf	Rather tolerant
Metribuzin pre-emergence	Rather susceptible
Metribuzin post-emergence	Rather susceptible

Storability

Dormancy* (2=short; 8=long)	5,4 
Pressure spots	
Fusarium	9 
Frying/crisping colour in October*	7,3 
Frying/crisping colour in February*	6,8 
Frying/crisping colour in May*	5,7 

*Potatoes stored at 8°C without sprout inhibitors

Marketable yield

Bruising index (0-50)	23 
Washability	6 
Common scab	7 
Powdery scab	6 
% IRS	2 
% Hollow heart	3 
Mechanical Damage	Rather susceptible

LADY JANE



Planting

Be aware of the risk of internals on sandy soils.
Wait for eyes to open before planting.
Plant deep and make big ridges in order to avoid tuber greening.

Size	75 cm	90 cm	Plant/ha
28/35	28 cm	23 cm	47600
35/50	38 cm	32 cm	35100
50/60	44 cm	37 cm	30300
60 +	45 cm	38 cm	29600
50/60 CUT	35 cm	29 cm	38100
60 + CUT	36 cm	30 cm	37000



Fertiliser input (kg/ha):

Depending on the soil supply and applicable national legislation; total amount required by the variety.



Nitrogen:	(N)	200	50	250
Phosphate:	(P ₂ O ₅)	110		110
Potassium:	(K ₂ O)	250	100	350

To prevent mineral deficiencies, apply these elements in time (B, Mg, Mn, Ca)

Apply potassium just before planting and during the growing season.
Despite a very strong foliage, splitting of nitrogen supply is needed.



Crop protection

Rather susceptible to the use of 450 g/ha of Metribuzin at pre-and post-emergence.
Manage late blight control as usual : spray when there is risk, with full dose and at normal frequency.



Harvest and storage

The tuber bulking takes place during foliar senescence.
Harvest carefully; susceptible to bruising.
Short dormancy; after harvest and wound healing cool down quickly, keep the temperature stable and start with sprout control.
Ethylene in storage can influence the frying quality of this variety.



Storage:

7 °C
8 months

Key advice to succeed this variety:

Plant deep and make big ridges in order to avoid tuber greening.
Despite a very strong foliage, splitting of nitrogen supply is needed.
Apply potassium just before planting and during the growing season.
Harvest carefully; susceptible to bruising.

Harvest in
good conditions
to avoid
bruising

