

# JAZZY (0-40 mm)

FRANCELINE x CUPIDO



Building on potential

## Added value of the variety

Excellent taste, texture and cooking quality.  
 Very nice skin quality.  
 Variety with a very high tuber set.  
 Variety suitable for baby potato production.  
 Low susceptibility to common scab.



## Characteristics

Shape of tuber	L	
Av. Tuber length (Ø 50-60 mm)	9,3 cm	
Maturity (9=early, 4=late)	7,6	
Colour of skin	LY	
Colour of flesh	7,1	
Suitability for cooking	8 A	

## Soil adaptability

	Sandy	Light	Heavy
Yield (Relative)	55	67	67
DM (%)	17,5	19,2	19,3
UWW	316	350	352
Tuber size (Ø mm)	36	36	40

Ro1	3	
Ro2/3		
Ro5		
Pa2		
Pa3		
wz 1 (D1)	10	
wz 2 (G1)	4	
wz 6 (O1)	4	
wz 18 (T1)	6	
TRV	7	
Moptop virus	Tolerant	

## Climate adaptability

Drought tolerance	Susceptible
Secondary growth	7

## Fertiliser requirement

	N	P	K	Mg	B	Ca	Mn
Quantity	Average	High (+25%)	Low (-25%)	High (+25%)	Average	Average	Average
No. applications	1	1	1	4	1	1	1

## Chemical requirement

Foliar late blight	3
Tuber late blight	Susceptible
Early blight	Rather susceptible
Y-virus	6
Yntn-virus	7
Leafroll virus	
Black dot	Tolerant
Silver scurf	Tolerant
Metribuzin pre-emergence	Tolerant
Metribuzin post-emergence	Susceptible

## Storability

Dormancy* (2=short; 8=long)	2,0	
Pressure spots	Rather susceptible	
Fusarium	4	
Frying/crisping colour in October*	4,2	
Frying/crisping colour in February*	3,3	
Frying/crisping colour in May*	3,3	

\*Potatoes stored at 8°C without sprout inhibitors

## Marketable yield

Bruising index (0-50)	5
Washability	8
Common scab	7
Powdery scab	6
% IRS	0
% Hollow heart	0
Mechanical Damage	Rather tolerant

# JAZZY (0-40 mm)



## Planting


Before planting keep the seed quiet; plant them out of cold storage.  
Plant at a normal depth.  
Growing in bed system gives a real plus to optimise baby production results.  
Ask your sales representative for further advice  
Treat the seeds against rhizoctonia and silver scurf.

Size	75 cm	90 cm	Plant/ha
25/30	15 cm	13 cm	88900
30/40	20 cm	17 cm	66700
40/50	26 cm	22 cm	51300
30/50	00 cm	00 cm	
50/55	32 cm	27 cm	41700



## Fertiliser input (kg/ha):

Depending on the soil supply and applicable national legislation; total amount required by the variety.



Nitrogen:	(N)	150	--
Phosphate:	(P2O5)	160	150
Potassium:	(K2O)	250	160
			250

To prevent mineral deficiencies, apply these elements in time ( B, Mg, Mn, Ca )

In order to increase the tuber set, Phosphate application in the ridge is advised.



## Crop protection

Susceptible to the use of 150 g/ha of Metribuzin at post-emergence.  
Variety susceptible to foliar late blight, do not neglect blight control.  
Quick tuber growing at the end of the cycle, kill in time.



## Harvest and storage

Harvest as soon as skin set is done, approx. 18 days after haulm killing.  
Short dormancy; after harvest and wound healing cool down quickly,  
keep the temperature stable and start with sprout control.  
Start sprouting inhibitor application as soon as potatoes are dry.



### Storage:

2 °C  
6 months

### Key advice to succeed this variety:

Before planting keep the seed quiet; plant them out of cold storage.  
Growing in bed system gives a real plus to optimise baby production results.  
In order to increase the tuber set, Phosphate application in the ridge is advised.  
Harvest as soon as skin set is done, approx. 18 days after haulm killing.

Keep  
storage  
temperature  
stable

