LADY CHRISTL



WS73-3-391 x MANSOUR



Added value of the variety

 ${\it Excellent\ taste,\ texture\ and\ cooking\ quality.}$

Very early variety.

Variety with a very high tuber set.

Uniform tuber size with high marketable yield

Can be harvest directly from green field.



Characteristics

Shape of tuber	0	
Av. Tuber length (Ø 50-60 mm)	7,9 cm	
Maturity (9=early, 4=late)	8,0	
Colour of skin	Υ	
Colour of flesh	6,8	
Suitability for cooking	8 A	
		A B C D

Soil adaptability

	Sandy	Light	Heavy
Yield (Relative)	81	70	72
DM (%)	17,7	19,6	19,6
UWW	320	357	357
Tuber size (Ø mm)	46	49	54

Ro1	9	
Ro2/3	\circ	
Ro5	\circ	
Pa2	\circ	
Pa3	0	
wz 1 (D1)	10	
wz 2 (G1)	\circ	
wz 6 (O1)	7	
wz 18 (T1)	0	
TRV	0	
Moptop virus	Tolerant	

Climate adaptability

Drought tolerance		
Secondary growth	8	

Chemical requirement

Foliar late blight	3 🔘	
Tuber late blight	Susceptible	
Early blight	Tolerant	
Y-virus	9 🔵	
Yntn-virus	10 🔵	
Leafroll virus	Susceptible	
Black dot	Susceptible	
Silver scurf	Susceptible	
Metribuzin pre-emergence	Rather susceptible	
Metribuzin post-emergence	Rather susceptible	

Storability

Dormancy* (2=short; 8=long)	3,4	
Pressure spots		
Fusarium		6 🔵
Frying/crisping colour in October*		4,4 🔘
Frying/crisping colour in February*		3,2 🔘
Frying/crisping colour in May*		3,0
*Potatoes stored at 8°C without sprout inhibitors		

Marketable yield

Bruising index (0-50)	14 🔵
Washability	7
Common scab	8 🔵
Powdery scab	3 🔘
% IRS	13 🔘
% Hollow heart	0
Mechanical Damage	Rather tolerant

Fertiliser requirement

	N	P	K	Mg	В	Ca	Mn
Quantity	Low (-25%)	Average	Average	Average	Average	Average	Average
No. applications	1	1	1	6	1	1	1

LADY CHRISTL





Planting

Only presprout in suitable chitting trays.


Plant in warm soils in order to avoid rhizoctonia.

Size	75 cm	90 cm	Plant/ha
28/30	18 cm	15 cm	74100
30/40	24 cm	20 cm	55600
40/50	30 cm	25 cm	44400
50/60	36 cm	30 cm	37000



Fertiliser input (kg/ha):

Depending on the soil supply and applicable national legislation; total amount required by the variety.



To prevent mineral deficiencies, apply these elements in time (B, Mg, Mn, Ca)

Be aware of Magnesium (Mg) deficiency.



Crop protection

Slightly susceptible to Metribuzin.

Variety susceptible to foliar late blight, do not neglect blight control.



Harvest and storage

Can be harvest directly from green field. Variety not suitable for long term storage.



Storage:

4 °C

2 months

Key advice to succeed this variety:

Plant in warm soils in order to avoid rhizoctonia. Be aware of Magnesium (Mg) deficiency. Slightly susceptible to Metribuzin. Can be harvest directly from green field.













